

Remote Measurements of Stratospheric Composition by Balloon-borne FTIR Solar Absorption Spectrometry.

G.C.Toon, J.-F.Blavier, B.Sen, and J.T.Szeto,
(Jet Propulsion Laboratory, MS 183-301, California
Institute of Technology, 48(10 Oak Grove Drive,
Pasadena, California, 91109; 818-354-8259)

The Mk IV Interferometer, a high resolution $1\text{--}11\text{ cm}^{-1}$ spectrometer built at JPL, for remote atmospheric measurements, has made 5 successful mid-latitude (35°N) balloon flights since 1989. Operated in solar absorption mode, each measured spectrum covered the entire $650\text{--}5450\text{ cm}^{-1}$ spectral region at 0.01 cm^{-1} resolution. Least squares fitting of computed spectra to those observed was employed to quantify over 20 different atmospheric gases simultaneously including ICN , Cl_2 , N_2O , OCS , CH_2Cl_2 , CF_2Cl_2 , CHCl_3 , CCl_4 , O_3 , NO , NO_2 , HNO_3 , N_2O_5 , ClONO_2 , HOCl , HCl , HF , COF_2 , CO , C_2H_6 , H_2O , HDO , N_2 , O_2 and CO_2 . The data reduction process will be described and the major sources of error will be examined. In particular, the last three named gases, having well known atmospheric abundances, are important in establishing the true observation geometry and temperature profile, which otherwise are major sources of uncertainty. Data taken during balloon ascent, sunset, sunrise and balloon descent have been separately inverted to yield sets of volume mixing ratio profiles over the 15-45 km altitude range. These not only provide diurnal information on the short-lived species, but also test the consistency of the other long-lived gases.

1. 1993 Fall Meeting
2. 010143674 (AGU number)
3. (a) G. C. Toon
MS 183-301
Jet Propulsion Laboratory
California Institute of Technology
4800 Oak Grove Drive
Pasadena, CA 91109

(t) Tel: 818-354 -8259

(c) Fax: 818-354-5148
4. A
5. (b) 0340 Middle Atm. Chem. Comp.
6. n/a
7. 5% SPIE Conference Berlin, 1992
8. \$50 (check enclosed)
9. C
10. Schedule before B.Sen
11. No